Cambridge IGCSE

Cambridge Assessment International Education
Cambridge International General Certificate of Secondary Education

CANDIDATE NAME									
CENTRE NUMBER						CANDIDATE NUMBER				

ADDITIONAL MATHEMATICS **0606/22**

Paper 2 **May/June 2019**

2 hours

Candidates answer on the Question Paper.

Additional Materials: Electronic calculator

READ THESE INSTRUCTIONS FIRST

Write your centre number, candidate number and name on all the work you hand in.
Write in dark blue or black pen.
You may use an HB pencil for any diagrams or graphs.
Do not use staples, paper clips, glue or correction fluid.
DO **NOT** WRITE IN ANY BARCODES.

Answer **all** the questions.
Give non-exact numerical answers correct to 3 significant figures, or 1 decimal place in the case of angles in degrees, unless a different level of accuracy is specified in the question.
The use of an electronic calculator is expected, where appropriate.
You are reminded of the need for clear presentation in your answers.

At the end of the examination, fasten all your work securely together.
The number of marks is given in brackets [ ] at the end of each question or part question.
The total number of marks for this paper is 80.

This document consists of **16** printed pages.

© UCLES 2019

Cambridge Assessment International Education

Mathematical Formulae

1. ALGEBRA

Quadratic Equation

For the equation $ax^2 + bx + c = 0$,

$$x = \frac{-b \pm \sqrt{b^2 - 4ac}}{2a}$$

Binomial Theorem

$$(a + b)^n = a^n + \binom{n}{1} a^{n-1}\, b + \binom{n}{2} a^{n-2}\, b^2 + \ldots + \binom{n}{r} a^{n-r}\, b^r + \ldots + b^n,$$

where n is a positive integer and $\binom{n}{r} = \frac{n!}{(n-r)!r!}$

2. TRIGONOMETRY

Identities

$$\sin^2 A + \cos^2 A = 1$$

$$\sec^2 A = 1 + \tan^2 A$$

$$\operatorname{cosec}^2 A = 1 + \cot^2 A$$

Formulae for ΔABC

$$\frac{a}{\sin A} = \frac{b}{\sin B} = \frac{c}{\sin C}$$

$$a^2 = b^2 + c^2 - 2bc \cos A$$

$$\Delta = \frac{1}{2} bc \sin A$$

© UCLES 2019

1 Given that $y = \dfrac{\sin x}{\ln x^2}$, find an expression for $\dfrac{\mathrm{d}y}{\mathrm{d}x}$. [4]

View solution

2 Find the values of k for which the equation $(k-1)x^2 + kx - k = 0$ has real and distinct roots. [4]

View solution

© UCLES 2019

3 **(i)** Given that $x-2$ is a factor of ax^3-12x^2+5x+6, use the factor theorem to show that $a=4$. [2]

View solution

(ii) Showing all your working, factorise $4x^3-12x^2+5x+6$ and hence solve $4x^3-12x^2+5x+6=0$. [4]

View solution

© UCLES 2019

4 A circle has diameter x which is increasing at a constant rate of $0.01\,\text{cm}\,\text{s}^{-1}$. Find the exact rate of change of the area of the circle when $x = 6\,\text{cm}$. [5]

View solution

5 **(i)** Express $5x^2 - 15x + 1$ in the form $p(x+q)^2 + r$, where p, q and r are constants. [3]

View solution

(ii) Hence state the least value of $x^2 - 3x + 0.2$ and the value of x at which this occurs. [2]

View solution

© UCLES 2019

6 **(a)** State the order of the matrix $\begin{pmatrix} 0 & 1 & 4 & 8 \\ 5 & 8 & 1 & 6 \end{pmatrix}$. [1]

View solution

(b) $\mathbf{A} = \begin{pmatrix} 2 & -4 \\ -1 & 3 \end{pmatrix}$

View solution

(i) Find $\mathbf{A}^{-1}$. [2]

(ii) Hence, given that $\mathbf{ABA} = \mathbf{I}$, find the matrix $\mathbf{B}$. [3]

View solution

© UCLES 2019

7 **(a)** Solve $\lg(x^2-3)=0$. [2]

View solution

(b) **(i)** Show that, for $a>0$, $\dfrac{\ln a^{\sin(2x+5)}+\ln\left(\dfrac{1}{a}\right)}{\ln a}$ may be written as $\sin(2x+5)+k$, where k is an integer. [3]

View solution

(ii) Hence find $\displaystyle\int \frac{\ln a^{\sin(2x+5)}+\ln\left(\frac{1}{a}\right)}{\ln a}\,\mathrm{d}x$. [3]

View solution

© UCLES 2019

8 **(a)** In the binomial expansion of $\left(a-\frac{x}{2}\right)^6$, the coefficient of x^3 is 120 times the coefficient of x^5. Find the possible values of the constant a. [4]

View solution

(b) **(i)** Expand $(1+2x)^{20}$ in ascending powers of x, as far as the term in x^3. Simplify each term. [2]

View solution

(ii) Use your expansion to show that the value of 0.98^{20} is 0.67 to 2 decimal places. [2]

View solution

© UCLES 2019

9 **(a)** Solve $6\sin^2 x - 13\cos x = 1$ for $0° \leqslant x \leqslant 360°$. [4]

View solution

(b) **(i)** Show that, for $-\frac{\pi}{2} < y < \frac{\pi}{2}$, $\frac{4\tan y}{\sqrt{1+\tan^2 y}}$ can be written in the form $a\sin y$, where a is an integer. [3]

View solution

(ii) Hence solve $\frac{4\tan y}{\sqrt{1+\tan^2 y}} + 3 = 0$ for $-\frac{\pi}{2} < y < \frac{\pi}{2}$ radians. [1]

View solution

10 **(a)** Find the unit vector in the direction of $5\mathbf{i} - 15\mathbf{j}$. [2]

View solution

(b) The position vectors of points A and B relative to an origin O are $\begin{pmatrix} 3 \\ -5 \end{pmatrix}$ and $\begin{pmatrix} 12 \\ 7 \end{pmatrix}$ respectively. The point C lies on AB such that $AC : CB$ is $2 : 1$.

(i) Find the position vector of C relative to O. [3]

View solution

The point D lies on OB such that $OD : OB$ is $1 : \lambda$ and $\overrightarrow{DC} = \begin{pmatrix} 6 \\ 1.25 \end{pmatrix}$.

(ii) Find the value of λ. [3]

View solution

11 The velocity, v m s^{-1}, of a particle travelling in a straight line, t seconds after passing through a fixed point O, is given by $v = \dfrac{4}{(t+1)^3}$.

(i) Explain why the direction of motion of the particle never changes. [1]

View solution

(ii) Showing all your working, find the acceleration of the particle when $t = 5$. [3]

View solution

(iii) Find an expression for the displacement of the particle from O after t seconds. [3]

View solution

(iv) Find the distance travelled by the particle in the fourth second. [2]

View solution

© UCLES 2019

12 **(a)** The functions f and g are defined by

$$\mathrm{f}(x) = 5x - 2 \quad \text{for } x > 1,$$
$$\mathrm{g}(x) = 4x^2 - 9 \quad \text{for } x > 0.$$

View solution

(i) State the range of g. [1]

(ii) Find the domain of gf. [1]

View solution

(iii) Showing all your working, find the exact solution of $\mathrm{gf}(x) = 45$. [3]

View solution

Question 12(b) is printed on the next page.

(b) The function h is defined by $\mathrm{h}(x) = \sqrt{x^2 - 1}$ for $x \leqslant -1$.

(i) State the geometrical relationship between the graphs of $y = \mathrm{h}(x)$ and $y = \mathrm{h}^{-1}(x)$. [1]

View solution

(ii) Find an expression for $\mathrm{h}^{-1}(x)$. [3]

View solution

Permission to reproduce items where third-party owned material protected by copyright is included has been sought and cleared where possible. Every reasonable effort has been made by the publisher (UCLES) to trace copyright holders, but if any items requiring clearance have unwittingly been included, the publisher will be pleased to make amends at the earliest possible opportunity.

To avoid the issue of disclosure of answer-related information to candidates, all copyright acknowledgements are reproduced online in the Cambridge Assessment International Education Copyright Acknowledgements Booklet. This is produced for each series of examinations and is freely available to download at www.cambridgeinternational.org after the live examination series.

Cambridge Assessment International Education is part of the Cambridge Assessment Group. Cambridge Assessment is the brand name of the University of Cambridge Local Examinations Syndicate (UCLES), which itself is a department of the University of Cambridge.

© UCLES 2019