Cambridge Assessment International Education

Cambridge IGCSE™

CANDIDATE NAME									
CENTRE NUMBER						CANDIDATE NUMBER				

ADDITIONAL MATHEMATICS **0606/11**

Paper 1 **May/June 2022**

2 hours

You must answer on the question paper.

No additional materials are needed.

INSTRUCTIONS

- Answer **all** questions.
- Use a black or dark blue pen. You may use an HB pencil for any diagrams or graphs.
- Write your name, centre number and candidate number in the boxes at the top of the page.
- Write your answer to each question in the space provided.
- Do **not** use an erasable pen or correction fluid.
- Do **not** write on any bar codes.
- You should use a calculator where appropriate.
- You must show all necessary working clearly; no marks will be given for unsupported answers from a calculator.
- Give non-exact numerical answers correct to 3 significant figures, or 1 decimal place for angles in degrees, unless a different level of accuracy is specified in the question.

INFORMATION

- The total mark for this paper is 80.
- The number of marks for each question or part question is shown in brackets [ ].

This document has **16** pages. Any blank pages are indicated.

© UCLES 2022

Mathematical Formulae

1. ALGEBRA

Quadratic Equation

For the equation $ax^2 + bx + c = 0$,

$$x = \frac{-b \pm \sqrt{b^2 - 4ac}}{2a}$$

Binomial Theorem

$$(a+b)^n = a^n + \binom{n}{1}a^{n-1}b + \binom{n}{2}a^{n-2}b^2 + \ldots + \binom{n}{r}a^{n-r}b^r + \ldots + b^n$$

where n is a positive integer and $\binom{n}{r} = \frac{n!}{(n-r)!r!}$

Arithmetic series $u_n = a + (n-1)d$

$$S_n = \frac{1}{2}n(a+l) = \frac{1}{2}n\{2a + (n-1)d\}$$

Geometric series $u_n = ar^{n-1}$

$$S_n = \frac{a(1-r^n)}{1-r} \quad (r \neq 1)$$

$$S_\infty = \frac{a}{1-r} \quad (|r| < 1)$$

2. TRIGONOMETRY

Identities

$$\sin^2 A + \cos^2 A = 1$$
$$\sec^2 A = 1 + \tan^2 A$$
$$\operatorname{cosec}^2 A = 1 + \cot^2 A$$

Formulae for ΔABC

$$\frac{a}{\sin A} = \frac{b}{\sin B} = \frac{c}{\sin C}$$

$$a^2 = b^2 + c^2 - 2bc\cos A$$

$$\Delta = \frac{1}{2}bc\sin A$$

© UCLES 2022

1 Find constants a, b and c such that $\dfrac{\sqrt{p}\,q^{\frac{2}{3}}r^{-3}}{\left(pq^{-1}\right)^2 r^{-1}} = p^a q^b r^c$. [3]

View solution

2 A particle moves in a straight line such that its displacement, s metres, from a fixed point, at time t seconds, $t \geqslant 0$, is given by $s = \left(1 + 3t\right)^{-\frac{1}{2}}$.

(a) Find the exact speed of the particle when $t = 1$. [3]

View solution

(b) Show that the acceleration of the particle will never be zero. [2]

View solution

3 A function f is such that $\mathrm{f}(x) = \ln(2x+1)$, for $x > -\frac{1}{2}$.

(a) Write down the range of f. [1]

View solution

A function g is such that $\mathrm{g}(x) = 5x - 7$, for $x \in \mathbb{R}$.

(b) Find the exact solution of the equation $\mathrm{gf}(x) = 13$. [3]

View solution

(c) Find the solution of the equation $\mathrm{f}'(x) = \mathrm{g}^{-1}(x)$. [6]

View solution

4 (a)



The diagram shows the graph of $y = |f(x)|$, where $f(x)$ is a cubic. Find the possible expressions for $f(x)$. [3]

(b) (i) On the axes below, sketch the graph of $y = |2x + 1|$ and the graph of $y = |4(x - 1)|$, stating the coordinates of the points where the graphs meet the coordinate axes. [3]



(ii) Find the exact solutions of the equation $|2x+1| = |4(x-1)|$. [4]

View solution

5 **(a)** Find the vector which is in the opposite direction to $\begin{pmatrix} 15 \\ -8 \end{pmatrix}$ and has a magnitude of 8.5. [2]

View solution

(b) Find the values of a and b such that $5\begin{pmatrix} 3a \\ b \end{pmatrix} + \begin{pmatrix} 2a+1 \\ 2 \end{pmatrix} = 6\begin{pmatrix} b+a \\ 2 \end{pmatrix}$. [3]

View solution

6 **(a)** Write down the values of k for which the line $y = k$ is a tangent to the curve $y = 4\sin\left(x + \frac{\pi}{4}\right) + 10$. [2]

View solution

(b) (i) Show that $\frac{1+\tan\theta}{1-\cos\theta}+\frac{1-\tan\theta}{1+\cos\theta}=\frac{2(1+\sin\theta)}{\sin^2\theta}$. [4]

View solution

(ii) Hence solve the equation $\frac{1+\tan\theta}{1-\cos\theta}+\frac{1-\tan\theta}{1+\cos\theta}=3$, for $0° \leqslant \theta \leqslant 360°$. [4]

View solution

© UCLES 2022

7 **(a)** The first three terms of an arithmetic progression are $\lg 3$, $3\lg 3$, $5\lg 3$. Given that the sum to n terms of this progression can be written as $256\lg 81$, find the value of n. [5]

View solution

(b) DO NOT USE A CALCULATOR IN THIS PART OF THE QUESTION.

The first three terms of a geometric progression are $\ln 256$, $\ln 16$, $\ln 4$. Find the sum to infinity of this progression, giving your answer in the form $p \ln 2$. [4]

View solution

8 DO NOT USE A CALCULATOR IN THIS QUESTION.

(a) Find the exact coordinates of the points of intersection of the curve $y = x^2 + 2\sqrt{5}x - 20$ and the line $y = 3\sqrt{5}x + 10$. [4]



View solution

(b) It is given that $\tan\theta = \dfrac{\sqrt{3}-1}{2+\sqrt{3}}$, for $0 < \theta < \dfrac{\pi}{2}$. Find $\text{cosec}^2\theta$ in the form $a + b\sqrt{3}$, where a and b are constants. [5]

View solution

9 A circle, centre O and radius r cm, has a sector OAB of fixed area $10\,\text{cm}^2$. Angle AOB is θ radians and the perimeter of the sector is P cm.

(a) Find an expression for P in terms of r. [3]

View solution

(b) Find the value of r for which P has a stationary value. [3]

View solution

(c) Determine the nature of this stationary value. [2]

View solution

(d) Find the value of θ at this stationary value. [1]

View solution

© UCLES 2022

10 The normal to the curve $y = \tan\left(3x + \frac{\pi}{2}\right)$ at the point P with coordinates $(p, -1)$, where $0 < p \leqslant \frac{\pi}{6}$, meets the x-axis at the point A and the y-axis at the point B. Find the exact coordinates of the mid-point of AB. [10]

View solution

BLANK PAGE

Permission to reproduce items where third-party owned material protected by copyright is included has been sought and cleared where possible. Every reasonable effort has been made by the publisher (UCLES) to trace copyright holders, but if any items requiring clearance have unwittingly been included, the publisher will be pleased to make amends at the earliest possible opportunity.

To avoid the issue of disclosure of answer-related information to candidates, all copyright acknowledgements are reproduced online in the Cambridge Assessment International Education Copyright Acknowledgements Booklet. This is produced for each series of examinations and is freely available to download at www.cambridgeinternational.org after the live examination series.

Cambridge Assessment International Education is part of Cambridge Assessment. Cambridge Assessment is the brand name of the University of Cambridge Local Examinations Syndicate (UCLES), which is a department of the University of Cambridge.

© UCLES 2022