Cambridge Assessment International Education

Cambridge IGCSE™

CANDIDATE NAME									
CENTRE NUMBER						CANDIDATE NUMBER				

ADDITIONAL MATHEMATICS **0606/21**

Paper 2 **May/June 2022**

2 hours

You must answer on the question paper.

No additional materials are needed.

INSTRUCTIONS

- Answer **all** questions.
- Use a black or dark blue pen. You may use an HB pencil for any diagrams or graphs.
- Write your name, centre number and candidate number in the boxes at the top of the page.
- Write your answer to each question in the space provided.
- Do **not** use an erasable pen or correction fluid.
- Do **not** write on any bar codes.
- You should use a calculator where appropriate.
- You must show all necessary working clearly; no marks will be given for unsupported answers from a calculator.
- Give non-exact numerical answers correct to 3 significant figures, or 1 decimal place for angles in degrees, unless a different level of accuracy is specified in the question.

INFORMATION

- The total mark for this paper is 80.
- The number of marks for each question or part question is shown in brackets [ ].

© UCLES 2022

Mathematical Formulae

1. ALGEBRA

Quadratic Equation

For the equation $ax^2 + bx + c = 0$,

$$x = \frac{-b \pm \sqrt{b^2 - 4ac}}{2a}$$

Binomial Theorem

$$(a+b)^n = a^n + \binom{n}{1}a^{n-1}b + \binom{n}{2}a^{n-2}b^2 + \ldots + \binom{n}{r}a^{n-r}b^r + \ldots + b^n$$

where n is a positive integer and $\binom{n}{r} = \frac{n!}{(n-r)!r!}$

Arithmetic series $u_n = a + (n-1)d$

$$S_n = \frac{1}{2}n(a+l) = \frac{1}{2}n\{2a + (n-1)d\}$$

Geometric series $u_n = ar^{n-1}$

$$S_n = \frac{a(1-r^n)}{1-r} \quad (r \neq 1)$$

$$S_\infty = \frac{a}{1-r} \quad (|r| < 1)$$

2. TRIGONOMETRY

Identities

$$\sin^2 A + \cos^2 A = 1$$
$$\sec^2 A = 1 + \tan^2 A$$
$$\operatorname{cosec}^2 A = 1 + \cot^2 A$$

Formulae for ΔABC

$$\frac{a}{\sin A} = \frac{b}{\sin B} = \frac{c}{\sin C}$$

$$a^2 = b^2 + c^2 - 2bc\cos A$$

$$\Delta = \frac{1}{2}bc\sin A$$

© UCLES 2022

1 **(a)** Solve the equation $5^{w-1} = 12$, giving your answer correct to 2 decimal places. [2]

View solution

(b) Solve the equation $x^{\frac{2}{3}} - 5x^{\frac{1}{3}} + 6 = 0$. [3]

View solution

2 **(a)** Write $2\lg x - \left(\lg(x+6) + \lg 3\right)$ as a single logarithm to base 10. [2]

View solution

(b) Hence solve the equation $2\lg x - \left(\lg(x+6) + \lg 3\right) = 0$. [4]

View solution

© UCLES 2022

3 Variables x and y are such that when $\sqrt[3]{y}$ is plotted against x^2, a straight line passing through the points (9, 8) and (16, 1) is obtained. Find y as a function of x. [4]

View solution

4 The polynomial $\mathrm{p}(x) = mx^3 - 17x^2 + nx + 6$ has a factor $x - 3$. It has a remainder of -12 when divided by $x + 1$. Find the remainder when $\mathrm{p}(x)$ is divided by $x - 2$. [6]

View solution

© UCLES 2022

5 **(a)** **(i)** Write down, in ascending powers of x, the first three terms in the expansion of $(1+4x)^n$. Simplify each term. [2]

View solution

(ii) In the expansion of $(1+4x)^n(1-4x)$ the coefficient of x^2 is 6032. Given that $n > 0$, find the value of n. [3]

View solution

(b) Find the term independent of x in the expansion of $\left(\frac{x}{2}-\frac{8}{x^4}\right)^{10}$. [2]

View solution

© UCLES 2022

6 **(a)** **(i)** A 5-digit number is to be formed from the seven digits 0, 1, 2, 3, 4, 5, 6. Each digit can be used at most once in any number and the number does not start with 0. Find the number of ways in which this can be done. [2]

View solution

(ii) Find how many of these 5-digit numbers are even. [3]

View solution

(b) A team of 7 people is to be selected from a group of 9 women and 6 men. Find the number of different teams that can be selected which include at least one man. [2]

View solution

(c) **(i)** Show that ${}^nC_3 + {}^nC_2 = \frac{1}{6}(n^3 - n)$ for $n \geqslant 3$. [5]

View solution

(ii) Hence solve the equation ${}^nC_3 + {}^nC_2 = 4n$ where $n \geqslant 3$. [2]

View solution

© UCLES 2022

7 Variables x and y are such that $y = \dfrac{(1+\sin 3x)^4}{\sqrt{x}}$. Use differentiation to find the approximate change in y when x increases from 1.9 to $1.9+h$, where h is small. [6]

View solution

8 In this question, **i** is a unit vector due east and **j** is a unit vector due north. Distances are measured in kilometres and time is measured in hours.

At 09 00, ship A leaves a point P with position vector $5\mathbf{i}+16\mathbf{j}$ relative to an origin O. It sails with a constant speed of $6\sqrt{3}$ on a bearing of 120°.

(a) Show that the velocity vector of A is $9\mathbf{i}-3\sqrt{3}\,\mathbf{j}$. [2]

View solution

(b) Find the position vector of A at 12 00. [1]

View solution

(c) At 11 00 ship B leaves a point Q with position vector $29\mathbf{i}+16\mathbf{j}$. It sails with constant velocity $-12\sqrt{3}\,\mathbf{j}$. Write down the position vector of B, t hours after it starts sailing. [1]

View solution

(d) Find the distance between the two ships at 12 00. [3]

View solution

© UCLES 2022

9 In this question all lengths are in metres.



The diagram shows a water container in the shape of a triangular prism. The depth of water in the container is h. The container has length 5. The water in the container forms a prism with a uniform cross-section that is an equilateral triangle of side x.

(a) Show that the volume, V, of the water is given by $V = \frac{5\sqrt{3}h^2}{3}$. [4]

View solution

(b) Water is pumped into the container at a rate of $0.5\ \text{m}^3$ per minute. Find the rate at which the depth of the water is increasing when the depth of the water is 0.1 m. [4]

View solution

10 **(a)** Differentiate $x\ln x - 2x$ with respect to x. Simplify your answer. [2]

View solution

(b) A curve is such that $\frac{d^2y}{dx^2} = \left(\frac{x+1}{\sqrt{x}}\right)^2$. It is given that $\frac{dy}{dx} = \frac{e^2}{2} + 2e$ at the point $\left(e, \frac{e^3}{6} + e^2\right)$.
Using your answer to **part (a)**, find the exact equation of the curve. [8]

View solution

Question 11 is printed on the next page.

© UCLES 2022

11

View solution



The diagram shows part of the curves $y = e^{\frac{x}{2}}$ and $y = \cos 5x$ and part of the line $x = \frac{\pi}{4}$. The curves intersect at A. The curve $y = \cos 5x$ cuts the x-axis at B. The line $x = \frac{\pi}{4}$ cuts the x-axis at C and the curve $y = e^{\frac{x}{2}}$ at D. Find the exact area of the shaded region, $ABCD$. [7]

Permission to reproduce items where third-party owned material protected by copyright is included has been sought and cleared where possible. Every reasonable effort has been made by the publisher (UCLES) to trace copyright holders, but if any items requiring clearance have unwittingly been included, the publisher will be pleased to make amends at the earliest possible opportunity.

To avoid the issue of disclosure of answer-related information to candidates, all copyright acknowledgements are reproduced online in the Cambridge Assessment International Education Copyright Acknowledgements Booklet. This is produced for each series of examinations and is freely available to download at www.cambridgeinternational.org after the live examination series.

Cambridge Assessment International Education is part of Cambridge Assessment. Cambridge Assessment is the brand name of the University of Cambridge Local Examinations Syndicate (UCLES), which is a department of the University of Cambridge.

© UCLES 2022