Cambridge Assessment International Education

Cambridge IGCSE™

CANDIDATE NAME									
CENTRE NUMBER						CANDIDATE NUMBER				

ADDITIONAL MATHEMATICS **0606/11**

Paper 1 **May/June 2024**

2 hours

You must answer on the question paper.

No additional materials are needed.

INSTRUCTIONS

- Answer **all** questions.
- Use a black or dark blue pen. You may use an HB pencil for any diagrams or graphs.
- Write your name, centre number and candidate number in the boxes at the top of the page.
- Write your answer to each question in the space provided.
- Do **not** use an erasable pen or correction fluid.
- Do **not** write on any bar codes.
- You should use a calculator where appropriate.
- You must show all necessary working clearly; no marks will be given for unsupported answers from a calculator.
- Give non-exact numerical answers correct to 3 significant figures, or 1 decimal place for angles in degrees, unless a different level of accuracy is specified in the question.

INFORMATION

- The total mark for this paper is 80.
- The number of marks for each question or part question is shown in brackets [ ].

© UCLES 2024

Mathematical Formulae

1. ALGEBRA

Quadratic Equation

For the equation $ax^2 + bx + c = 0$,

$$x = \frac{-b \pm \sqrt{b^2 - 4ac}}{2a}$$

Binomial Theorem

$$(a+b)^n = a^n + \binom{n}{1}a^{n-1}b + \binom{n}{2}a^{n-2}b^2 + \ldots + \binom{n}{r}a^{n-r}b^r + \ldots + b^n$$

where n is a positive integer and $\binom{n}{r} = \frac{n!}{(n-r)!r!}$

Arithmetic series $u_n = a + (n-1)d$

$$S_n = \frac{1}{2}n(a+l) = \frac{1}{2}n\{2a + (n-1)d\}$$

Geometric series $u_n = ar^{n-1}$

$$S_n = \frac{a(1-r^n)}{1-r} \quad (r \neq 1)$$

$$S_\infty = \frac{a}{1-r} \quad (|r| < 1)$$

2. TRIGONOMETRY

Identities

$$\sin^2 A + \cos^2 A = 1$$
$$\sec^2 A = 1 + \tan^2 A$$
$$\operatorname{cosec}^2 A = 1 + \cot^2 A$$

Formulae for ΔABC

$$\frac{a}{\sin A} = \frac{b}{\sin B} = \frac{c}{\sin C}$$

$$a^2 = b^2 + c^2 - 2bc\cos A$$

$$\Delta = \frac{1}{2}bc\sin A$$

© UCLES 2024

1 **(a)** On the axes, sketch the graph of $y = -\frac{1}{5}(x+2)(2x-1)(x+5)$, stating the intercepts with the axes. [3]

View solution



(b) Hence solve the inequality $-\frac{1}{5}(x+2)(2x-1)(x+5) \geqslant 0$. [2]

View solution

2 DO NOT USE A CALCULATOR IN THIS QUESTION.

The polynomial p is such that $\mathrm{p}(x) = 6x^3 - 35x^2 + 34x + 45$.

(a) Find $\mathrm{p}(x)$ in the form $(2x-5)\mathrm{q}(x) + r$, where $\mathrm{q}(x)$ is a polynomial and r is a constant. [3]

View solution

(b) Hence write the expression $\mathrm{p}(x) - 5$ as a product of linear factors. [2]

View solution

(c) Hence write down the solutions of the equation $\mathrm{p}(x) = 5$. [1]

View solution

3 **(a)** Write $1+\lg(x^2-1)-2\lg(x-1)$, where $x>1$, as a single logarithm to base 10. Give your answer in its simplest form. [4]

View solution

(b) Solve the equation $4\log_5(x+1)=9\log_{(x+1)}5$, giving your answers in the form $a+b\sqrt{c}$, where a, b and c are constants. [5]

View solution

© UCLES 2024

4 **(a)** The first three terms, in ascending powers of x, in the expansion of $(3+px)^n$ are $243+810x+qx^2$, where n, p and q are constants. Find the values of n, p and q. [5]

View solution

(b) Find the term independent of y in the expansion of $\left(2y-\frac{1}{3y^2}\right)^6$. Give your answer in exact form. [2]

View solution

© UCLES 2024

5 **(a)** The diagram shows the graph of $y = a\cos bx + c$, for $-360° \leqslant x \leqslant 360°$, where a, b and c are constants. Find the values of a, b and c. [3]

View solution



(b) The line $y = p$ is a tangent to the curve $y = 3 - 2\sin 6\theta$. Write down the possible values of p. [2]

View solution

© UCLES 2024

6 Find $\int_2^4 \left(\frac{2}{2x-3} - \frac{3}{(3x-5)^2} \right) \mathrm{d}x$, giving your answer in exact form. [4]

View solution

7 Given that $2 + \cot\theta = 3x$ and $\sin\theta = \sqrt{y}$, find y in terms of x. [3]

View solution

© UCLES 2024

8 Solve the equation $4\sin^2\left(2\alpha - \frac{\pi}{3}\right) = 1$ for $-\frac{\pi}{2} \leqslant \alpha \leqslant \frac{\pi}{2}$. Give your answers in terms of π. [5]

View solution

9 **(a)** Solve the following simultaneous equations.

View solution

$$e^{x+y} \times e^{3x-2y} = 1$$

$$x^2y = 256$$

[5]

(b) Solve the equation $10e^{(2x-1)} - 11 = 6e^{(1-2x)}$, giving your answer in exact form. [4]

View solution

10 In this question, all distances are in metres and time, t, is in seconds.

A particle P is at a fixed point O at time $t = 0$.
The velocity, v, of P is given by $v = 3\sin 2t$ for $t \geqslant 0$.

(a) Find the exact value of t for which the velocity is zero for the first time after P leaves O. [2]

View solution

(b) Find an expression, in terms of t, for the displacement of P from O at time t. [4]

View solution

(c) Find the distance travelled by P for $0 \leqslant t \leqslant \pi$. [3]

View solution

11 The tangent to the curve $y = (3x-1)^{\frac{1}{3}}$ at the point where $x = 3$ meets the coordinate axes at the points A and B. The point with coordinates (a, a) lies on the perpendicular bisector of the line AB. Find the exact value of a. [10]

View solution

Continuation of working space for Question 11.

Question 12 is printed on the next page.

12 **(a)** It is given that $y = \dfrac{\ln 3x}{x^2}$ for $x > 0$.

Find $\dfrac{\mathrm{d}y}{\mathrm{d}x}$. Give your answer in the form $\dfrac{A + B\ln 3x}{x^3}$, where A and B are integers. [4]

View solution

(b) Hence find $\displaystyle\int \frac{\ln 3x}{x^3}\,\mathrm{d}x$. [4]

View solution

Permission to reproduce items where third-party owned material protected by copyright is included has been sought and cleared where possible. Every reasonable effort has been made by the publisher (UCLES) to trace copyright holders, but if any items requiring clearance have unwittingly been included, the publisher will be pleased to make amends at the earliest possible opportunity.

To avoid the issue of disclosure of answer-related information to candidates, all copyright acknowledgements are reproduced online in the Cambridge Assessment International Education Copyright Acknowledgements Booklet. This is produced for each series of examinations and is freely available to download at www.cambridgeinternational.org after the live examination series.

Cambridge Assessment International Education is part of Cambridge Assessment. Cambridge Assessment is the brand name of the University of Cambridge Local Examinations Syndicate (UCLES), which is a department of the University of Cambridge.

© UCLES 2024