Cambridge IGCSE

Cambridge International Examinations
Cambridge International General Certificate of Secondary Education

CANDIDATE NAME									
CENTRE NUMBER						CANDIDATE NUMBER				

ADDITIONAL MATHEMATICS **0606/23**

Paper 2 **October/November 2015**

2 hours

Candidates answer on the Question Paper.

Additional Materials: Electronic calculator

READ THESE INSTRUCTIONS FIRST

Write your Centre number, candidate number and name on all the work you hand in.
Write in dark blue or black pen.
You may use an HB pencil for any diagrams or graphs.
Do not use staples, paper clips, glue or correction fluid.
DO **NOT** WRITE IN ANY BARCODES.

Answer **all** the questions.
Give non-exact numerical answers correct to 3 significant figures, or 1 decimal place in the case of angles in degrees, unless a different level of accuracy is specified in the question.
The use of an electronic calculator is expected, where appropriate.
You are reminded of the need for clear presentation in your answers.

At the end of the examination, fasten all your work securely together.
The number of marks is given in brackets [ ] at the end of each question or part question.
The total number of marks for this paper is 80.

This document consists of **16** printed pages.

© UCLES 2015

CAMBRIDGE International Examinations

Mathematical Formulae

1. ALGEBRA

Quadratic Equation

For the equation $ax^2 + bx + c = 0$,

$$x = \frac{-b \pm \sqrt{b^2 - 4ac}}{2a}$$

Binomial Theorem

$$(a + b)^n = a^n + \binom{n}{1} a^{n-1} b + \binom{n}{2} a^{n-2} b^2 + \ldots + \binom{n}{r} a^{n-r} b^r + \ldots + b^n,$$

where n is a positive integer and $\binom{n}{r} = \frac{n!}{(n-r)!r!}$

2. TRIGONOMETRY

Identities

$$\sin^2 A + \cos^2 A = 1$$

$$\sec^2 A = 1 + \tan^2 A$$

$$\operatorname{cosec}^2 A = 1 + \cot^2 A$$

Formulae for ΔABC

$$\frac{a}{\sin A} = \frac{b}{\sin B} = \frac{c}{\sin C}$$

$$a^2 = b^2 + c^2 - 2bc \cos A$$

$$\Delta = \frac{1}{2} bc \sin A$$

© UCLES 2015

1 Find the equation of the tangent to the curve $y = x^3 + 3x^2 - 5x - 7$ at the point where $x = 2$. [5]

View solution

© UCLES 2015

2 Find the values of k for which the line $y = 2x + k + 2$ cuts the curve $y = 2x^2 + (k + 2)x + 8$ in two distinct points. [6]

View solution

3 **(a)** Given that $y = \dfrac{x^3}{2 - x^2}$, find $\dfrac{\mathrm{d}y}{\mathrm{d}x}$. [3]

View solution

(b) Given that $y = x\sqrt{4x + 6}$, show that $\dfrac{\mathrm{d}y}{\mathrm{d}x} = \dfrac{k(x + 1)}{\sqrt{4x + 6}}$ and state the value of k. [3]

View solution

© UCLES 2015

4 Solve the following simultaneous equations, giving your answers for both x and y in the form $a + b\sqrt{3}$, where a and b are integers.

View solution

$$2x + y = 9$$

$$\sqrt{3}x + 2y = 5$$ [5]

5 The roots of the equation $x^3 + ax^2 + bx + c = 0$ are 1, 3 and 3. Show that $c = -9$ and find the value of a and of b. [4]

View solution

6 Solve the following equation.

$$\log_2(29x - 15) = 3 + \frac{2}{\log_x 2}$$ [5]

View solution

7 The velocity, v ms^{-1}, of a particle travelling in a straight line, t seconds after passing through a fixed point O, is given by $v = \dfrac{10}{(2+t)^2}$.

(i) Find the acceleration of the particle when $t = 3$. [3]

View solution

(ii) Explain why the particle never comes to rest. [1]

View solution

(iii) Find an expression for the displacement of the particle from O after time t s. [3]

View solution

(iv) Find the distance travelled by the particle between $t = 3$ and $t = 8$. [2]

View solution

© UCLES 2015

8 **(i)** Prove that $\sec^2 x + \operatorname{cosec}^2 x = \sec^2 x \operatorname{cosec}^2 x$. [4]

View solution

(ii) Hence, or otherwise, solve $\sec^2 x + \operatorname{cosec}^2 x = 4\tan^2 x$ for $90° < x < 270°$. [4]

View solution

9 Given that $f(x) = 3x^2 + 12x + 2$,

(i) find values of a, b and c such that $f(x) = a(x + b)^2 + c$, [3]

View solution

(ii) state the minimum value of $f(x)$ and the value of x at which it occurs, [2]

View solution

(iii) solve $f\left(\frac{1}{y}\right) = 0$, giving each answer for y correct to 2 decimal places. [3]

View solution

© UCLES 2015

10 **(i)** Given that $\frac{\mathrm{d}}{\mathrm{d}x}(\mathrm{e}^{2-x^2}) = kx\mathrm{e}^{2-x^2}$, state the value of k. [1]

View solution

(ii) Using your result from part **(i)**, find $\int 3x\mathrm{e}^{2-x^2}\mathrm{d}x$. [2]

View solution

(iii) Hence find the area enclosed by the curve $y = 3x\mathrm{e}^{2-x^2}$, the x-axis and the lines $x = 1$ and $x = \sqrt{2}$. [2]

View solution

© UCLES 2015

(iv) Find the coordinates of the stationary points on the curve $y = 3xe^{2-x^2}$. [4]

View solution

11 The trees in a certain forest are dying because of an unknown virus.

The number of trees, N, surviving t years after the onset of the virus is shown in the table below.

t	1	2	3	4	5	6
N	2000	1300	890	590	395	260

The relationship between N and t is thought to be of the form $N = Ab^{-t}$.

(i) Transform this relationship into straight line form. [1]

View solution

(ii) Using the given data, draw this straight line on the grid below. [3]

View solution



(iii) Use your graph to estimate the value of A and of b. [3]

View solution

If the trees continue to die in the same way, find

(iv) the number of trees surviving after 10 years, [1]

View solution

(v) the number of years taken until there are only 10 trees surviving. [2]

View solution

Question 12 is printed on the next page.

© UCLES 2015

12 A plane that can travel at 250 kmh^{-1} in still air sets off on a bearing of 070°. A wind with speed $w\,\text{kmh}^{-1}$ from the south blows the plane off course so that the plane actually travels on a bearing of 060°.

Find, in kmh^{-1}, the resultant speed V of the plane and the windspeed w. [5]

View solution

Permission to reproduce items where third-party owned material protected by copyright is included has been sought and cleared where possible. Every reasonable effort has been made by the publisher (UCLES) to trace copyright holders, but if any items requiring clearance have unwittingly been included, the publisher will be pleased to make amends at the earliest possible opportunity.

To avoid the issue of disclosure of answer-related information to candidates, all copyright acknowledgements are reproduced online in the Cambridge International Examinations Copyright Acknowledgements Booklet. This is produced for each series of examinations and is freely available to download at www.cie.org.uk after the live examination series.

Cambridge International Examinations is part of the Cambridge Assessment Group. Cambridge Assessment is the brand name of University of Cambridge Local Examinations Syndicate (UCLES), which is itself a department of the University of Cambridge.

© UCLES 2015