Cambridge Assessment International Education

Cambridge IGCSE™

CANDIDATE NAME									
CENTRE NUMBER						CANDIDATE NUMBER				

ADDITIONAL MATHEMATICS **0606/22**

Paper 2 **October/November 2023**

2 hours

You must answer on the question paper.

No additional materials are needed.

INSTRUCTIONS

- Answer **all** questions.
- Use a black or dark blue pen. You may use an HB pencil for any diagrams or graphs.
- Write your name, centre number and candidate number in the boxes at the top of the page.
- Write your answer to each question in the space provided.
- Do **not** use an erasable pen or correction fluid.
- Do **not** write on any bar codes.
- You should use a calculator where appropriate.
- You must show all necessary working clearly; no marks will be given for unsupported answers from a calculator.
- Give non-exact numerical answers correct to 3 significant figures, or 1 decimal place for angles in degrees, unless a different level of accuracy is specified in the question.

INFORMATION

- The total mark for this paper is 80.
- The number of marks for each question or part question is shown in brackets [ ].

© UCLES 2023

Mathematical Formulae

1. ALGEBRA

Quadratic Equation

For the equation $ax^2 + bx + c = 0$,

$$x = \frac{-b \pm \sqrt{b^2 - 4ac}}{2a}$$

Binomial Theorem

$$(a+b)^n = a^n + \binom{n}{1}a^{n-1}b + \binom{n}{2}a^{n-2}b^2 + \ldots + \binom{n}{r}a^{n-r}b^r + \ldots + b^n$$

where n is a positive integer and $\binom{n}{r} = \frac{n!}{(n-r)!r!}$

Arithmetic series $u_n = a + (n-1)d$

$$S_n = \frac{1}{2}n(a+l) = \frac{1}{2}n\{2a + (n-1)d\}$$

Geometric series $u_n = ar^{n-1}$

$$S_n = \frac{a(1-r^n)}{1-r} \quad (r \neq 1)$$

$$S_\infty = \frac{a}{1-r} \quad (|r| < 1)$$

2. TRIGONOMETRY

Identities

$$\sin^2 A + \cos^2 A = 1$$
$$\sec^2 A = 1 + \tan^2 A$$
$$\operatorname{cosec}^2 A = 1 + \cot^2 A$$

Formulae for ΔABC

$$\frac{a}{\sin A} = \frac{b}{\sin B} = \frac{c}{\sin C}$$

$$a^2 = b^2 + c^2 - 2bc\cos A$$

$$\Delta = \frac{1}{2}bc\sin A$$

© UCLES 2023

1 **(a)** A straight line passes through the points (4, 23) and (−8, 29). Find the point of intersection, P, of this line with the line $y = 2x + 5$. [5]

View solution

(b) Find the distance of P from the origin. [2]

View solution

© UCLES 2023

2 Find the non-zero value of k for which the line $y = -2x - 6k - 1$ is a tangent to the curve $y = x(x + 2k)$. [5]

View solution

3 DO NOT USE A CALCULATOR IN THIS QUESTION.

A cylinder has base radius $(2+\sqrt{3})$ m and volume $\pi(16+9\sqrt{3})\text{m}^3$. Find the exact value of its height, giving your answer in its simplest form. [4]

View solution

4 Solve the following equations.

(a) $\dfrac{(e^{x+1})^2}{\sqrt{e^x}} = 10$ [4]

View solution

(b) $2\log_9 y - \log_9(4y-9) = \dfrac{1}{2}$ [5]

View solution

5 **(a)** Find the equation of the normal to the curve $y = x^3 - 7x^2 + 12x - 5$ at the point (1, 1). [5]

View solution

(b) Find the x-coordinates of the two points where the normal cuts the curve again. Give your answers in the form $x = a \pm \sqrt{b}$ where a and b are integers. [5]

View solution

© UCLES 2023

6 Find the exact value of $\int_2^3 \frac{(x+2)^2}{x}\,\mathrm{d}x$. [6]

View solution

7 A particle is travelling in a straight line. Its displacement, s metres, from the origin at time t seconds is given by $s = 1.5e^{2t} + 2e^{-2t} - t$.

(a) Find expressions for the velocity, $v\,\text{ms}^{-1}$, and acceleration, $a\,\text{ms}^{-2}$, of the particle. [3]

View solution

(b) Find the time, T seconds, when the particle is at rest. [4]

View solution

(c) Find the acceleration of the particle at time T seconds. [2]

View solution

© UCLES 2023

8 A curve has equation $y = x \sin 2x$.

(a) Find $\frac{\mathrm{d}y}{\mathrm{d}x}$. [2]

View solution

(b) Find the equation of the tangent to the curve at $x = \frac{\pi}{4}$. [3]

View solution

(c) Use your answer to **part (a)** to find the exact value of $\int_0^{\frac{\pi}{6}} 2x\cos 2x\mathrm{d}x$. [5]

View solution

9 **(a)** An arithmetic progression has twelve terms. The sum of the first three terms is -36 and the sum of the last three terms is 72. Find the first term and the common difference. [5]

View solution

(b) The first three terms of a geometric progression are 1, 1.2 and 1.44. Find the smallest value of n such that the sum of the first n terms is greater than 500. [5]

View solution

© UCLES 2023

10 **(a)** By writing $\cot x$ and $\tan x$ in terms of $\cos x$ and $\sin x$, show that

$$\frac{\sin x}{1-\cot x}+\frac{\cos x}{1-\tan x}=\sin x+\cos x.$$ [5]

View solution

(b) Solve the equation $9\cot x + 3\operatorname{cosec} x = \tan x$, for $0° < x < 360°$. [5]

View solution

BLANK PAGE

Permission to reproduce items where third-party owned material protected by copyright is included has been sought and cleared where possible. Every reasonable effort has been made by the publisher (UCLES) to trace copyright holders, but if any items requiring clearance have unwittingly been included, the publisher will be pleased to make amends at the earliest possible opportunity.

To avoid the issue of disclosure of answer-related information to candidates, all copyright acknowledgements are reproduced online in the Cambridge Assessment International Education Copyright Acknowledgements Booklet. This is produced for each series of examinations and is freely available to download at www.cambridgeinternational.org after the live examination series.

Cambridge Assessment International Education is part of Cambridge Assessment. Cambridge Assessment is the brand name of the University of Cambridge Local Examinations Syndicate (UCLES), which is a department of the University of Cambridge.

© UCLES 2023