Cambridge Assessment International Education

# Cambridge IGCSE™

| | | | | | | | | | |
|---|---|---|---|---|---|---|---|---|---|
| CANDIDATE NAME | | | | | | | | | |
| CENTRE NUMBER | | | | | | CANDIDATE NUMBER | | | | |

**CHEMISTRY** **0620/52**

Paper 5 Practical Test **October/November 2021**

**1 hour 15 minutes**

You must answer on the question paper.

You will need: The materials and apparatus listed in the confidential instructions

**INSTRUCTIONS**

- Answer **all** questions.
- Use a black or dark blue pen. You may use an HB pencil for any diagrams or graphs.
- Write your name, centre number and candidate number in the boxes at the top of the page.
- Write your answer to each question in the space provided.
- Do **not** use an erasable pen or correction fluid.
- Do **not** write on any bar codes.
- You may use a calculator.
- You should show all your working and use appropriate units.

**INFORMATION**

- The total mark for this paper is 40.
- The number of marks for each question or part question is shown in brackets [ ].
- Notes for use in qualitative analysis are provided in the question paper.

| For Examiner's Use | |
|---|---|
| 1 | |
| 2 | |
| 3 | |
| Total | |

This document has **12** pages. Any blank pages are indicated.

© UCLES 2021

**1** You are going to investigate the reaction between two different solutions of aqueous sodium carbonate, labelled solution **K** and solution **L**, and two different solutions of dilute hydrochloric acid, labelled acid **M** and acid **N**.

**Read all of the instructions carefully before starting the experiments.**

**Instructions**
You are going to do three experiments.

View solution

**(a)** *Experiment 1*

- Fill the burette with solution **K**. Run some of solution **K** out of the burette so that the level of solution **K** is on the burette scale.
- Use the measuring cylinder to pour 25 $cm^3$ of acid **M** into the conical flask.
- Add five drops of methyl orange indicator to the conical flask.
- Stand the conical flask on a white tile.
- Slowly add solution **K** from the burette to the conical flask, while swirling the flask, until the solution just changes colour.
- Record the burette readings in the table and complete the table.

| | Experiment 1 |
|---|---|
| final burette reading / $cm^3$ | |
| initial burette reading / $cm^3$ | |
| volume of solution **K** added / $cm^3$ | |

*Experiment 2*

- Empty the conical flask and rinse it with distilled water.
- Refill the burette with solution **K**. Run some of solution **K** out of the burette so that the level of solution **K** is on the burette scale.
- Use the measuring cylinder to pour 25 $cm^3$ of acid **N** into the conical flask.
- Add five drops of methyl orange indicator to the conical flask.
- Stand the conical flask on a white tile.
- Slowly add solution **K** from the burette to the conical flask, while swirling the flask, until the solution just changes colour.
- Record the burette readings in the table and complete the table.

| | Experiment 2 |
|---|---|
| final burette reading / $cm^3$ | |
| initial burette reading / $cm^3$ | |
| volume of solution **K** added / $cm^3$ | |

© UCLES 2021

*Experiment 3*

- Empty the burette and rinse it with distilled water.
- Empty the conical flask and rinse it with distilled water.
- Fill the burette with solution **L**. Run some of solution **L** out of the burette so that the level of solution **L** is on the burette scale.
- Use the measuring cylinder to pour 25 $cm^3$ of acid **N** into the conical flask.
- Add five drops of methyl orange indicator to the conical flask.
- Stand the conical flask on a white tile.
- Slowly add solution **L** from the burette to the conical flask, while swirling the flask, until the solution just changes colour.
- Record the burette readings in the table and complete the table.

| | Experiment 3 |
|---|---|
| final burette reading / $cm^3$ | |
| initial burette reading / $cm^3$ | |
| volume of solution **L** added / $cm^3$ | |

[5]

**(b)** State the colour change observed at the end-point in the conical flask in Experiment 1.




from ..... to .....

[1]

**(c)** Describe one **other** observation made when solution **K** is added to acid **M** in Experiment 1.

..... [1]

**(d)** **(i)** Compare the volumes of solution **K** used in Experiment 1 and Experiment 2.




.....

..... [2]

**(ii)** Suggest why different volumes of solution **K** were needed in Experiment 1 and Experiment 2.

..... [1]

**(e)** Deduce the volume of solution **L** required to reach the end-point if Experiment 3 is repeated using acid **M** in place of acid **N**.




volume of solution **L** = ..... $cm^3$ [1]

**(f)** Explain why the conical flask was rinsed with water at the start of Experiment 2 and Experiment 3.

..... [1]

**(g)** At the start of Experiment 3 the burette was rinsed with water.

View solution

Describe an additional step that should have been done after rinsing the burette with water but before filling the burette with solution **L**. Explain your answer.

.....

..... [2]

**(h)** Explain why the conical flask is placed on a white tile.

View solution

..... [1]

**(i)** Describe how the reliability of the results of the experiments can be confirmed.

View solution

.....

..... [1]

**(j)** State **one** source of error in Experiment 1. Suggest an improvement to reduce this error.

View solution

source of error .....

improvement .....

[2]

[Total: 18]

**2** You are provided with solid **O** and liquid **P**.
Do the following tests on the two substances, recording all of your observations at each stage.

**tests on solid O**

**(a)** Use a spatula to transfer approximately one third of solid **O** to a boiling tube.
Heat the boiling tube gently and then strongly.
Record your observations.

View solution

.....

..... [2]

Place the remaining solid **O** in a boiling tube. Add about 10 $cm^3$ of distilled water to the boiling tube.
Place a stopper in the boiling tube and shake the tube to dissolve solid **O** and form solution **O**.

Divide solution **O** into three approximately equal portions in one boiling tube and two test-tubes.

**(b)** To the first portion of solution **O**, in the boiling tube, add approximately 2 $cm^3$ of aqueous sodium hydroxide.

**Keep the product for part (c).**

View solution

Record your observations.

..... [1]

**(c)** Warm the product from **(b)**. Test any gas produced.
Record your observations.
Identify the gas produced.

View solution

observations .....

identity of gas .....
[2]

**(d)** To the second portion of solution **O** add about 1 cm depth of dilute nitric acid followed by a few drops of aqueous silver nitrate.
Record your observations.

View solution

..... [1]

**(e)** To the third portion of solution **O**, add the aqueous chlorine.
Record your observations.

View solution

..... [1]

**(f)** Identify solid **O**.

View solution

.....

..... [2]

**tests on liquid P**

**(g)** Add a few drops of liquid **P** to the test-tube containing aqueous bromine. Replace the stopper in the test-tube and shake the test-tube.
Record your observations.

View solution

.....

..... [2]

**(h)** Add a few drops of liquid **P** to a crucible. Apply a lighted splint to the surface of liquid **P** in the crucible.
Record your observations.

View solution

.....

..... [1]

**(i)** To the remaining liquid **P** add a few drops of aqueous alkaline potassium manganate(VII). Replace the stopper in the test-tube and shake the test-tube. Leave the test-tube to stand for a few minutes.
Record your observations.

View solution

..... [2]

**(j)** State what conclusions can be made about liquid **P**.

View solution

.....

..... [2]

[Total: 16]

**3** Cobalt is a metal. Cobalt is between copper and iron in the reactivity series. The mineral spherocobaltite contains the compound cobalt(II) carbonate and no other metal ions. Cobalt(II) carbonate is insoluble in water and reacts with dilute acids to form an aqueous solution of a salt.

Describe how you would obtain a sample of cobalt metal starting with a large lump of spherocobaltite. You have access to all normal laboratory apparatus and chemicals.

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

..... [6]

View solution

**BLANK PAGE**

**BLANK PAGE**

**BLANK PAGE**

**Notes for use in qualitative analysis**
**Tests for anions**

| anion | test | test result |
|---|---|---|
| carbonate ($CO_3^{2-}$) | add dilute acid | effervescence, carbon dioxide produced |
| chloride ($Cl^-$) [in solution] | acidify with dilute nitric acid, then add aqueous silver nitrate | white ppt. |
| bromide ($Br^-$) [in solution] | acidify with dilute nitric acid, then add aqueous silver nitrate | cream ppt. |
| iodide ($I^-$) [in solution] | acidify with dilute nitric acid, then add aqueous silver nitrate | yellow ppt. |
| nitrate ($NO_3^-$) [in solution] | add aqueous sodium hydroxide, then aluminium foil; warm carefully | ammonia produced |
| sulfate ($SO_4^{2-}$) [in solution] | acidify, then add aqueous barium nitrate | white ppt. |
| sulfite ($SO_3^{2-}$) | add dilute hydrochloric acid, warm gently and test for the presence of sulfur dioxide | sulfur dioxide produced will turn acidified aqueous potassium manganate(VII) from purple to colourless |

**Tests for aqueous cations**

| cation | effect of aqueous sodium hydroxide | effect of aqueous ammonia |
|---|---|---|
| aluminium ($Al^{3+}$) | white ppt., soluble in excess, giving a colourless solution | white ppt., insoluble in excess |
| ammonium ($NH_4^+$) | ammonia produced on warming | – |
| calcium ($Ca^{2+}$) | white ppt., insoluble in excess | no ppt., or very slight white ppt. |
| chromium(III) ($Cr^{3+}$) | green ppt., soluble in excess | grey-green ppt., insoluble in excess |
| copper(II) ($Cu^{2+}$) | light blue ppt., insoluble in excess | light blue ppt., soluble in excess, giving a dark blue solution |
| iron(II) ($Fe^{2+}$) | green ppt., insoluble in excess | green ppt., insoluble in excess |
| iron(III) ($Fe^{3+}$) | red-brown ppt., insoluble in excess | red-brown ppt., insoluble in excess |
| zinc ($Zn^{2+}$) | white ppt., soluble in excess, giving a colourless solution | white ppt., soluble in excess, giving a colourless solution |

© UCLES 2021

**Tests for gases**

| gas | test and test result |
|---|---|
| ammonia ($NH_3$) | turns damp red litmus paper blue |
| carbon dioxide ($CO_2$) | turns limewater milky |
| chlorine ($Cl_2$) | bleaches damp litmus paper |
| hydrogen ($H_2$) | 'pops' with a lighted splint |
| oxygen ($O_2$) | relights a glowing splint |
| sulfur dioxide ($SO_2$) | turns acidified aqueous potassium manganate(VII) from purple to colourless |

**Flame tests for metal ions**

| metal ion | flame colour |
|---|---|
| lithium ($Li^+$) | red |
| sodium ($Na^+$) | yellow |
| potassium ($K^+$) | lilac |
| copper(II) ($Cu^{2+}$) | blue-green |

Permission to reproduce items where third-party owned material protected by copyright is included has been sought and cleared where possible. Every reasonable effort has been made by the publisher (UCLES) to trace copyright holders, but if any items requiring clearance have unwittingly been included, the publisher will be pleased to make amends at the earliest possible opportunity.

To avoid the issue of disclosure of answer-related information to candidates, all copyright acknowledgements are reproduced online in the Cambridge Assessment International Education Copyright Acknowledgements Booklet. This is produced for each series of examinations and is freely available to download at www.cambridgeinternational.org after the live examination series.

Cambridge Assessment International Education is part of the Cambridge Assessment Group. Cambridge Assessment is the brand name of the University of Cambridge Local Examinations Syndicate (UCLES), which itself is a department of the University of Cambridge.